Taking Your Hepatitis C Therapy:

Harvoni® with or without Ribavirin

Ledipasvir/Sofosbuvir = Harvoni®

- ▶ Dose: 1 tablet (90/400 mg) daily, taken at the same time each day
- ▶ You can take with or without food



- ▶ Inform your provider of all other medicines, supplements or over-the-counter products you take, especially omeprazole, ranitidine, famotidine or any other acid-reducing medicines
- ▶ Store at room temperature (< 86°F) away from direct sunlight

Do You Need to Take Ribavirin? □Yes □No



- Dose: 200 mg capsules or tablets; your dose may change during the course of treatment
- ► Take with food to minimize stomach upset such as nausea and vomiting
- Do not consider pregnancy until at least 6 months after treatment for either partner has ended
- ► Store at room temperature (< 86°F) away from direct sunlight

My Daily Medication Schedule Start Date:	_ End Date: Expected Treatment Duration: weeks
Morning Time: am Take 1 Ledipasvir/Sofosbuvir (Harvoni®) table	t daily
Morning Time: am Take Ribavirin with food	Evening Time: pm

Notes: - Harvoni® and Ribavirin can be taken together or at different times, but keep to the same time of day for each drug.

- If you take acid-reducing medicines, take Harvoni® in the morning on an empty stomach together with your acid-reducing medicine. If you also take Ribavirin, take this later in the morning or at lunch with food.

If You Miss a Dose:

Ledipasvir/Sofosbuvir: Harvoni®

- ▶ If you miss a dose, TAKE THE MISSED DOSE THE SAME DAY as soon as you remember; take your next dose of Harvoni® at your regular time the next day
- ▶ Do not take more than 1 tablet of Harvoni® in a day
- ▶ DO NOT take two doses at the same time to make up for a missed dose
- Note: If you miss multiple doses, call your prescriber/clinic listed below

Ribavirin

▶ If you miss more than 1-2 days of Ribavirin, call your prescriber/clinic listed below

Prescriber/Clinic Contact:	or Refills:	
Prescriber/Clinic Contact:	Or Remis:	



