Taking Your Hepatitis C Therapy: Vosevi®

Sofosbuvir/Velpatasvir/Voxilaprevir = Vosevi

- ▶ Dose: 1 tablet (100/400 mg) daily, taken at the same time each day with food
- ▶ Inform your provider of all other medicines, supplements, or over-the-counter products you take, including St John's wart, amiodarone, seizure medications, or acid reducing medicine such as omeprazole
- ▶ Get tested for hepatitis B (HBV). Tell your provider if you are currently taking, or have ever taken, any medication for HBV.
- ▶ Store at room temperature (< 86°F) away from direct sunlight

Daily Medication Schedule Start Date: _____ End Date: _____

	Expected Treatment D	Ouration: week	s Test for Cure Date:
Morning Time: am Take 1 Sofosbuvir/Velpa	atasvir/Voxilaprevir (Vosev	vi™) tablet daily with	food.
Notes: If you take acid-reducing medicines, like omeprazole (Prilosec*), take Vosevi in the morning with food together with your acid-reducing medicine. If you need to take an antacid medicine that contains aluminum or magnesium, take it either 4 hours before or 4 hours after you take your dose of Vosevi			
If You Miss a Dose:			
Sofosbuvir/Velpatasvir/Voxilaprevir: V	osevi		
 If you miss a dose, TAKE THE MISSED DOSE THE SAME DAY as soon as you remember; take your next dose of Vosevi at your regular time the next day Do not take more than 1 tablet of Vosevi in a day DO NOT take two doses at the same time to make up for a missed dose Note: If you miss multiple doses, call your prescriber/clinic listed below 			
Prescriber/Clinic Contact:		For Refills:	

