

Taking Your Hepatitis C Therapy:

Mavyret®

Glecaprevir/Pibrentasvir = Mavyret®




- ▶ Dose: 3 tablets daily at the same time each day, taken with food
- ▶ Mavyret® comes in a bottle (28-day supply) or a 4-week carton. In a 4-week carton, each carton contains a 7-day supply of medications in blister packs; one blister pack = 3 pills per day)
- ▶ Tell your provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, herbs, and supplements
- ▶ Get tested for hepatitis B (HBV); tell your provider if you are currently taking, or have ever taken, any medication for HBV
- ▶ Store Mavyret® at room temperature (< 86° F)

Daily Medication Schedule

Start Date: _____ End Date: _____

Expected Treatment Duration: ____ weeks Test for Cure Date: _____

Time: ____ am/pm

Take 3 Mavyret® tablets daily with food. 

If You Miss a Dose:

Glecaprevir/Pibrentasvir = Mavyret®

- ▶ If you miss a dose (3 tablets) less than 18 hours from the usual time that Mavyret® should have been taken, then take the dose (3 tablets) as soon as possible, and then take the next dose at the usual time
- ▶ If you miss a dose (3 tablets) more than 18 hours from the usual time that Mavyret® should have been taken, then do not take that day's dose at all, and take the next dose (3 tablets) at the usual time
- ▶ Note: If you miss multiple doses, call your prescriber/clinic listed below

Prescriber/Clinic Contact: _____

For Refills: _____



U.S. Department of Veterans Affairs

Veterans Health Administration