

# Managing Side Effects of Mavyret™

(Glecaprevir/Pibrentasvir)

## Headache (13%)

- ▶ Maintain adequate fluid intake
- ▶ Try to get plenty of rest
- ▶ Take acetaminophen (Tylenol®)\* as advised by your provider; see instructions on page 2

## Fatigue (11%)

- ▶ Try low-impact exercise (walking or low-impact aerobics) as directed by your provider
- ▶ Maintain adequate fluid intake (at least 6 to 8 non-caffeinated 8-ounce glasses/day)
- ▶ Take a short nap during the day
- ▶ Lessen your work schedule if possible
- ▶ Eat well-balanced meals daily

## Nausea (8%)

- ▶ Eat smaller meals
- ▶ Avoid foods/smells that trigger nausea; eat cold foods
- ▶ Eat healthy foods; avoid greasy, spicy, acidic, or sweet foods
- ▶ Try ginger in ginger tea, ginger ale, or gingersnaps
- ▶ Eat some crackers or dry white toast if you feel sick in the morning
- ▶ Over-the-counter antacids or other medications may be recommended

## Diarrhea (3-7%)

- ▶ Eat fiber like bananas, white rice, applesauce, and white toast (the BRAT diet)
- ▶ Avoid foods that are spicy or acidic (like citrus)
- ▶ Avoid dairy products for 2-3 days after diarrhea resolves
- ▶ Maintain adequate fluid intake (at least 6-8 noncaffeinated 8-ounce glasses/day)
- ▶ Your provider may recommend loperamide (Imodium®)\* or methylcellulose (Citrucel®)\* or psyllium (Metamucil®)\*

\* Generic versions are usually less expensive than brand-name products. Brand names are provided as examples only, and their inclusion does not indicate an endorsement by the Veterans Health Administration or any other government agency. Also, if a particular brand name is not mentioned, it does not mean or imply that the product is unsatisfactory.



**U.S. Department  
of Veterans Affairs**

Veterans Health  
Administration

## Additional Instructions While on Treatment

In addition to taking your medication as prescribed, you will need to go to the laboratory for blood tests:

- ▶ within 4 weeks of starting treatment and as needed thereafter
- ▶ 12 weeks after you take your last dose of medication (to ensure you have achieved a cure)

You will be given specific dates to get your blood tested. It is important that you follow the schedule and discuss the results with your health care provider.

Notify your provider about all medications including over-the-counter products and dietary/herbal supplements that you are taking. Ask your provider before taking any new medications while on treatment.

Notify your provider of any adverse effects during treatment. Seek immediate medical attention if you develop any serious side effect such as dark urine or yellowing eyes or skin.

## Helpful Telephone Numbers

Telephone Advice Nurse: \_\_\_\_\_

Clinic: \_\_\_\_\_

Provider: \_\_\_\_\_

Glecaprevir/Pibrentasvir Support Line: 1-877-628-9738

## Other Medications Approved for Your Use if Checked Off by Your Provider

### For Aches/Pains

- Acetaminophen (Tylenol®): Take 1-2 tablets every 4-6 hours if needed, up to maximum of 2,000 mg a day. DO NOT TAKE MORE THAN 6 tablets a day of 325 mg OR 4 tablets a day of 500 mg acetaminophen (Tylenol®).
- Ibuprofen (Advil® or Motrin®) \_\_\_\_\_ mg: Take \_\_\_ tablet every 6 hours with food.
- Naproxen (Aleve®) \_\_\_\_\_ mg: Take \_\_\_ tablet twice a day with food.

### For Sleep

- Diphenhydramine (Benadryl®) \_\_\_\_\_ mg: Take \_\_\_ capsule/tablet at bedtime as needed.
- Trazodone \_\_\_\_\_ mg: Take \_\_\_ tablet at bedtime as needed.